

Lecture 4. Kernel Methods

Bao Wang

Department of Mathematics


Scientific Computing and Imaging Institute

University of Utah

Math 5750/6880, Fall 2023

Support Vector Machine

Assume the training data set is **linearly separable in feature space**, i.e. there exists at least one set of \mathbf{w} and b s.t. a function of the form $y(\mathbf{x}) = \mathbf{w}^\top \mathbf{x} + b$ satisfies $y(\mathbf{x}_n) > 0 (< 0)$ for points having $t_n = +1 (-1)$, i.e., $t_n y(\mathbf{x}_n) > 0$ for all training data points.



Which classifier is better? [Theoretical motivation: VC-dimension.]

The maximum margin solution is found by solving

$$\arg \max_{\mathbf{w}, b} \left\{ \frac{1}{\|\mathbf{w}\|} \min_n \left[t_n(\mathbf{w}^\top \phi(\mathbf{x}_n) + b) \right] \right\}. \quad (1)$$

Direct solution of this optimization problem would be very complex, and so we shall convert it into an equivalent problem that is much easier to solve.

For any κ , we have

$$\arg \max_{\mathbf{w}, b} \left\{ \frac{1}{\|\mathbf{w}\|} \min_n \left[t_n(\mathbf{w}^\top \phi(\mathbf{x}_n) + b) \right] \right\} = \arg \max_{\mathbf{w}, b} \left\{ \frac{1}{\|\kappa \mathbf{w}\|} \min_n \left[t_n(\kappa \mathbf{w}^\top \phi(\mathbf{x}_n) + \kappa b) \right] \right\}$$

Support Vector Machine

We can choose κ s.t. $\min_n [t_n(\kappa \mathbf{w}^\top \phi(\mathbf{x}_n) + \kappa b)] = 1$, and the problem becomes

$$\arg \max_{\mathbf{w}, b} \left\{ \frac{1}{\|\kappa \mathbf{w}\|} \right\},$$

s.t.

$$t_n(\kappa \mathbf{w}^\top \phi(\mathbf{x}_n) + \kappa b) \geq 1, \quad \forall n = 1, \dots, N.$$

The problem is equivalent to

$$\arg \max_{\mathbf{w}, b} \left\{ \frac{1}{\|\mathbf{w}\|} \right\},$$

s.t.

$$t_n(\mathbf{w}^\top \phi(\mathbf{x}_n) + b) \geq 1, \quad \forall n = 1, \dots, N.$$

Support Vector Machine: Dual Form

We introduce Lagrange multipliers $a_n \geq 0$, with one multiplier a_n for each of the constraints, giving the Lagrangian function¹

$$L(\mathbf{w}, b, \mathbf{a}) = \frac{1}{2} \|\mathbf{w}\|^2 - \sum_{n=1}^N a_n \{t_n(\mathbf{w}^\top \phi(\mathbf{x}_n) + b) - 1\}, \quad \mathbf{a} = (a_1, \dots, a_N)^\top. \quad (2)$$

Setting the derivatives of $L(\mathbf{w}, b, \mathbf{a})$ w.r.t. \mathbf{w} and b equal to zero, we have

$$\mathbf{w} = \sum_{n=1}^N a_n t_n \phi(\mathbf{x}_n); \quad 0 = \sum_{n=1}^N a_n t_n. \quad (3)$$

¹KKT condition: If we wish to minimize the function $f(\mathbf{x})$ s.t. $g(\mathbf{x}) \geq 0$, then we minimize the Lagrangian function $L(\mathbf{x}, \lambda) = f(\mathbf{x}) - \lambda g(\mathbf{x})$ w.r.t. \mathbf{x} , subject to $\lambda \geq 0$, $g(\mathbf{x}) \geq 0$, and $\lambda g(\mathbf{x}) = 0$.

Support Vector Machine: Dual Form

Eliminating \mathbf{w} and b from $L(\mathbf{w}, b, \mathbf{a})$ using these conditions then gives the *dual representation* of the maximum margin problem in which we maximize

$$\tilde{L}(\mathbf{a}) = \sum_{n=1}^N a_n - \frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N a_n a_m t_n t_m k(\mathbf{x}_n, \mathbf{x}_m), \quad k(\mathbf{x}_n, \mathbf{x}_m) = \phi(\mathbf{x}_n)^\top \phi(\mathbf{x}_m) \text{ is the kernel.} \quad (4)$$

with respect to \mathbf{a} subject to the constraints

$$a_n \geq 0; \quad \sum_{n=1}^N a_n t_n = 0. \quad (5)$$

The linear classifier becomes

$$y(\mathbf{x}) = \sum_{n=1}^N a_n t_n k(\mathbf{x}, \mathbf{x}_n) + b. \quad (6)$$

Support Vector Machine (Soft Margin)

$$\min_{\mathbf{w}, \xi_n} C \sum_{n=1}^N \xi_n + \frac{1}{2} \|\mathbf{w}\|^2, \quad C > 0, \quad (7)$$

s.t.

$$t_n(\mathbf{w}^\top \phi(\mathbf{x}_n) + b) \geq 1 - \xi_n, \quad \xi_n \geq 0.$$

Support Vector Machine (Soft Margin): Dual Form

The corresponding Lagrangian is given by

$$L(\mathbf{w}, b, \mathbf{a}) = \frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{n=1}^N \xi_n - \sum_{n=1}^N a_n \left\{ t_n y(\mathbf{x}_n) - 1 + \xi_n \right\} - \sum_{n=1}^N \mu_n \xi_n, \quad (8)$$

where $\{a_n \geq 0\}$ and $\{\mu_n \geq 0\}$ are Lagrange multipliers. The corresponding KKT conditions are

$$\begin{aligned} a_n &\geq 0 \\ t_n y(\mathbf{x}_n) - 1 + \xi_n &\geq 0 \\ a_n(t_n y(\mathbf{x}_n) - 1 + \xi_n) &= 0 \\ \mu_n &\geq 0 \\ \xi_n &\geq 0 \\ \mu_n \xi_n &= 0 \end{aligned} \quad (9)$$

where $n = 1, \dots, N$.

Support Vector Machine (Soft Margin): Dual Form

The dual Lagrangian in the form

$$\tilde{L}(\mathbf{a}) = \sum_{n=1}^N a_n - \frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N a_n a_m t_n t_m k(\mathbf{x}_n, \mathbf{x}_m), \quad (10)$$

Kernel and kernel trick

The dual form of SVM involves a kernel function $k(\mathbf{x}, \mathbf{x}')$.

Many linear parametric models can be re-cast into an equivalent 'dual representation' in which the prediction are based on linear combinations of a *kernel function* evaluated at the training data points. As we shall see, **for models which are based on a fixed nonlinear *feature space* mapping $\phi(\mathbf{x})$, the kernel function is given by the relation**

$$k(\mathbf{x}, \mathbf{x}') = \phi(\mathbf{x})^\top \phi(\mathbf{x}'). \quad (11)$$

Kernel trick: a.k.a. *kernel substitution*. If we have an algorithm formulated in such a way that the input vector \mathbf{x} enters only in the form of scalar products, then **we can replace that scalar product with some other choice of kernels.**

Dual representation of linear regression

Linear regression model: $y(\mathbf{x}) = \mathbf{w}^\top \phi(\mathbf{x})$.

Consider the following regularized sum-of-squares error function for linear regression

$$J(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{\mathbf{w}^\top \phi(\mathbf{x}_n) - t_n\}^2 + \frac{\lambda}{2} \mathbf{w}^\top \mathbf{w}, \quad \lambda \geq 0. \quad (12)$$

What is the dual representation of the linear regression model above?

Dual representation of linear regression

$$\nabla_{\mathbf{w}} J(\mathbf{w}) = 0 \Rightarrow \mathbf{w} = -\frac{1}{\lambda} \sum_{n=1}^N \{\mathbf{w}^\top \phi(\mathbf{x}_n) - t_n\} \phi(\mathbf{x}_n) = \sum_{n=1}^N a_n \phi(\mathbf{x}_n) = \Phi^\top \mathbf{a},$$

where Φ is the design matrix, whose n th row is given by $\phi(\mathbf{x}_n)^\top$. The vector $\mathbf{a} = (a_1, \dots, a_N)^\top$ with

$$a_n = -\frac{1}{\lambda} \{\mathbf{w}^\top \phi(\mathbf{x}_n) - t_n\}. \quad (13)$$

Dual representation of linear regression

We can reformulate the least squares algorithm in terms of the parameter vector \mathbf{a} instead of \mathbf{w} , resulting in a *dual representation*. Substitute $\mathbf{w} = \Phi^\top \mathbf{a}$ into $J(\mathbf{w})$ gives

$$J(\mathbf{a}) = \frac{1}{2} \mathbf{a}^\top \Phi \Phi^\top \Phi \Phi^\top \mathbf{a} - \mathbf{a}^\top \Phi \Phi^\top \mathbf{t} + \frac{1}{2} \mathbf{t}^\top \mathbf{t} + \frac{\lambda}{2} \mathbf{a}^\top \Phi \Phi^\top \mathbf{a}, \quad (14)$$

where $\mathbf{t} = (t_1, \dots, t_N)^\top$.

Dual representation of linear regression

Define the *Gram* matrix $\mathbf{K} = \Phi\Phi^\top \in \mathbb{R}^{N \times N}$ with $K_{nm} = \phi(\mathbf{x}_n)^\top \phi(\mathbf{x}_m) = k(\mathbf{x}_n, \mathbf{x}_m)$. In terms of the Gram matrix, the sum-of-squares error function can be written as

$$J(\mathbf{a}) = \frac{1}{2} \mathbf{a}^\top \mathbf{K} \mathbf{K} \mathbf{a} - \mathbf{a}^\top \mathbf{K} \mathbf{t} + \frac{1}{2} \mathbf{t}^\top \mathbf{t} + \frac{\lambda}{2} \mathbf{a}^\top \mathbf{K} \mathbf{a}. \quad (15)$$

$$\nabla_{\mathbf{a}} J(\mathbf{a}) = 0 \Rightarrow \mathbf{a} = (\mathbf{K} + \lambda \mathbf{I}_N)^{-1} \mathbf{t}.$$

Dual representation of linear regression

Substitute this back into the linear regression model, we obtain the following prediction for a new input \mathbf{x}

$$y(\mathbf{x}) = \mathbf{w}^\top \phi(\mathbf{x}) = \mathbf{a}^\top \Phi \phi(\mathbf{x}) = \mathbf{k}(\mathbf{x})^\top (\mathbf{K} + \lambda \mathbf{I}_N)^{-1} \mathbf{t}, \quad \text{Dual formulation!} \quad (16)$$

where we have defined the vector $\mathbf{k}(\mathbf{x})$ with elements $k_n(\mathbf{x}) = k(\mathbf{x}_n, \mathbf{x})$.

The dual formulation allows the solution to the least-squares problem to be expressed entirely in terms of the kernel function $k(\mathbf{x}, \mathbf{x}')$.

Note that the prediction at \mathbf{x} is given by a linear combination of the target values from the training set.

Dual representation of linear regression

Remark. In the dual formulation, \mathbf{a} is determined by inverting an $N \times N$ matrix, whereas in the original parameter space formulation we had to invert an $M \times M$ matrix to determine \mathbf{w} (How?). Note that typically $N \gg M$, **the dual formulation seems inferior to the original formulation.**

However, the advantage of the dual formulation is that it is expressed entirely in terms of the kernel function $k(\mathbf{x}, \mathbf{x}')$. We can therefore work in terms of kernels and avoid the explicit introducing the feature vector $\phi(\mathbf{x})$, allowing us to implicitly use feature spaces of high, even infinite, dimensionality.

Constructing kernels

Approach I. Choose a feature map $\phi(\mathbf{x}) \in \mathbb{R}^M$ and then use this to find the corresponding kernel:

$$k(\mathbf{x}, \mathbf{x}') = \phi(\mathbf{x})^\top \phi(\mathbf{x}') = \sum_{i=1}^M \phi_i(\mathbf{x}) \phi_i(\mathbf{x}') \quad (17)$$

where $\phi_i(x)$ are the basis functions.

Constructing kernels

Approach II. Direct construction: in this case, we must ensure that the function we choose is a valid kernel, i.e. it corresponds to a scalar product in some (perhaps infinite dimensional) feature space. As a simple example, consider a kernel function given by

$$k(\mathbf{x}, \mathbf{z}) = (\mathbf{x}^\top \mathbf{z})^2. \quad (18)$$

If we take the particular case of a 2D input space $\mathbf{x} = (x_1, x_2)$ we can expand out the terms and thereby identify the corresponding nonlinear feature mapping

$$k(\mathbf{x}, \mathbf{z}) = (\mathbf{x}^\top \mathbf{z})^2 = (x_1 z_1 + x_2 z_2)^2 = (x_1^2, \sqrt{2}x_1 x_2, x_2^2)(z_1^2, \sqrt{2}z_1 z_2, z_2^2)^\top = \phi(\mathbf{x})^\top \phi(\mathbf{z}). \quad (19)$$

We see that the feature mapping takes the form $\phi(\mathbf{x}) = (x_1^2, \sqrt{2}x_1 x_2, x_2^2)^\top$ and therefore comprises all possible second order terms, with a specific weighting between them.

Constructing kernels

When $k(\mathbf{x}, \mathbf{x}')$ represents a kernel function?

Lemma. *A symmetric function $k : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ implements an inner product in some Hilbert space if and only if it is positive semidefinite; namely, for all $\mathbf{x}_1, \dots, \mathbf{x}_m$, the Gram matrix, $G_{i,j} = k(\mathbf{x}_i, \mathbf{x}_j)$, is a positive semidefinite matrix.*

Constructing kernels

Proof. It is trivial to see that if k implements an inner product in some Hilbert space then the Gram matrix is positive semidefinite.

For the other direction, define the space of functions over \mathcal{X} as $\mathbb{R}^{\mathcal{X}} = \{f : \mathcal{X} \rightarrow \mathbb{R}\}$. For each $\mathbf{x} \in \mathcal{X}$ let $\psi(\mathbf{x})$ be the function $\mathbf{x} \rightarrow K(\cdot, \mathbf{x})$. Define a vector space by taking all linear combinations of elements of the form $K(\cdot, \mathbf{x})$. Define an inner product on this vector space to be

$$\left\langle \sum_i \alpha_i K(\cdot, \mathbf{x}_i), \sum_j \beta_j K(\cdot, \mathbf{x}'_j) \right\rangle = \sum_{i,j} \alpha_i \beta_j K(\mathbf{x}_i, \mathbf{x}'_j).$$

This is a valid inner product since it is symmetric (because K is symmetric), it is linear (immediate), and it is positive definite (it is easy to see that $K(\mathbf{x}, \mathbf{x}) \geq 0$ with equality only for $\psi(\mathbf{x})$ being the zero function). Clearly,

$$\langle \psi(\mathbf{x}), \psi(\mathbf{x}') \rangle = \langle K(\cdot, \mathbf{x}), K(\cdot, \mathbf{x}') \rangle = K(\mathbf{x}, \mathbf{x}'),$$

which concludes our proof.

Constructing kernels

Build them out of simpler kernels as building blocks.

Proposition. Given valid kernels $k_1(\mathbf{x}, \mathbf{x}')$ and $k_2(\mathbf{x}, \mathbf{x}')$, the following new kernels will also be valid:

$$k(\mathbf{x}, \mathbf{x}') = ck_1(\mathbf{x}, \mathbf{x}')$$

$$k(\mathbf{x}, \mathbf{x}') = f(\mathbf{x})k_1(\mathbf{x}, \mathbf{x}')f(\mathbf{x}')$$

$$k(\mathbf{x}, \mathbf{x}') = q(k_1(\mathbf{x}, \mathbf{x}'))$$

$$k(\mathbf{x}, \mathbf{x}') = \exp(k_1(\mathbf{x}, \mathbf{x}'))$$

$$k(\mathbf{x}, \mathbf{x}') = k_1(\mathbf{x}, \mathbf{x}') + k_2(\mathbf{x}, \mathbf{x}')$$

$$k(\mathbf{x}, \mathbf{x}') = k_1(\mathbf{x}, \mathbf{x}')k_2(\mathbf{x}, \mathbf{x}')$$

$$k(\mathbf{x}, \mathbf{x}') = k_3(\phi(\mathbf{x}), \phi(\mathbf{x}'))$$

$$k(\mathbf{x}, \mathbf{x}') = \mathbf{x}^\top \mathbf{A} \mathbf{x}'$$

$$k(\mathbf{x}, \mathbf{x}') = k_a(\mathbf{x}_a, \mathbf{x}'_a) + k_b(\mathbf{x}_b, \mathbf{x}'_b)$$

$$k(\mathbf{x}, \mathbf{x}') = k_a(\mathbf{x}_a, \mathbf{x}'_a)k_b(\mathbf{x}_b, \mathbf{x}'_b)$$

where $c > 0$ is a constant, $f(\cdot)$ is any function, $q(\cdot)$ is a polynomial with nonnegative coefficients, $\phi(\mathbf{x})$ is a function from \mathbf{x} to \mathbb{R}^M , $k_3(\cdot, \cdot)$ is a valid kernel in \mathbb{R}^M , \mathbf{A} is p.s.d, \mathbf{x}_a and \mathbf{x}_b are variables with $\mathbf{x} = (\mathbf{x}_a, \mathbf{x}_b)$, and k_a and k_b are valid kernel functions over their respective spaces.

Gaussian kernels

Let the original instance space be \mathbb{R} and consider the mapping ϕ where for each nonnegative integer $n \geq 0$ there exists an element $\phi(\mathbf{x})_n$ (n -th coordinate) that equals $\frac{1}{\sqrt{n!}} e^{-\frac{x^2}{2}} x^n$. Then,

$$\langle \phi(\mathbf{x}), \phi(\mathbf{x}') \rangle = \sum_{n=0}^{\infty} \left(\frac{1}{\sqrt{n!}} e^{-\frac{x^2}{2}} x^n \right) \left(\frac{1}{\sqrt{n!}} e^{-\frac{(x')^2}{2}} (x')^n \right) = e^{-\frac{x^2 + (x')^2}{2}} \sum_{n=0}^{\infty} \left(\frac{(xx')^n}{n!} \right) = e^{-\frac{\|x-x'\|^2}{2}}.$$

Here the feature space is of infinite dimension while evaluating the kernel is very simple. More generally, given a scalar $\sigma > 0$, the Gaussian kernel is defined to be

$$K(\mathbf{x}, \mathbf{x}') = e^{-\frac{\|\mathbf{x}-\mathbf{x}'\|^2}{2\sigma}}.$$

Intuitively, the Gaussian kernel sets the inner product in the feature space between \mathbf{x}, \mathbf{x}' to be close to zero if the instances are far away from each other (in the original domain) and close to 1 if they are close. σ is a parameter that controls the scale determining what we mean by "close".

The Gaussian kernel is also called the RBF kernel.

Polynomial kernels

The k degree polynomial kernel is defined to be

$$K(\mathbf{x}, \mathbf{x}') = (1 + \langle \mathbf{x}, \mathbf{x}' \rangle)^k.$$

We will show that there exists a mapping ϕ from the original space to some higher dimensional space for which $K(\mathbf{x}, \mathbf{x}') = \langle \phi(\mathbf{x}), \phi(\mathbf{x}') \rangle$. For simplicity, denote $x_0 = x'_0 = 1$. Then, we have

$$\begin{aligned} K(\mathbf{x}, \mathbf{x}') &= (1 + \langle \mathbf{x}, \mathbf{x}' \rangle)^k = \left(\sum_{j=0}^n x_j x'_j \right) \cdots \left(\sum_{j=0}^n x_j x'_j \right) = \sum_{J \in \{0,1,\dots,n\}^k} \prod_{i=1}^k x_{J_i} x'_{J_i} \\ &= \sum_{J \in \{0,1,\dots,n\}^k} \prod_{i=1}^k x_{J_i} \prod_{i=1}^k x'_{J_i}. \end{aligned}$$

Now, if we define $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^{(n+1)^k}$ such that for $J \in \{0,1,\dots,n\}^k$ there is an element of $\phi(\mathbf{x})$ that equals $\prod_{i=1}^k x_{J_i}$, we obtain that

$$K(\mathbf{x}, \mathbf{x}') = \langle \phi(\mathbf{x}), \phi(\mathbf{x}') \rangle.$$

Kernel trick

$$\langle \mathbf{x}, \mathbf{x}' \rangle \Rightarrow \langle \phi(\mathbf{x}), \phi(\mathbf{x}') \rangle \Rightarrow k(\mathbf{x}, \mathbf{x}').$$