



# Multigraded rings, diagonal subalgebras, and rational singularities

Kazuhiko Kurano <sup>a</sup>, Ei-ichi Sato <sup>b</sup>, Anurag K. Singh <sup>c,\*</sup>,  
Kei-ichi Watanabe <sup>d</sup>

<sup>a</sup> Department of Mathematics, Meiji University, Higashimita 1-1-1, Tama-ku, Kawasaki-shi 214-8571, Japan

<sup>b</sup> Department of Mathematics, Kyushu University, Hakozaki 6-10-1, Higashi-ku, Fukuoka-city 812-8581, Japan

<sup>c</sup> Department of Mathematics, University of Utah, 155 South 1400 East, Salt Lake City, UT 84112, USA

<sup>d</sup> Department of Mathematics, Nihon University, Sakura-Josui 3-25-40, Setagaya, Tokyo 156-8550, Japan

Received 27 March 2008

Available online 6 December 2008

Communicated by Luchezar L. Avramov

To Paul Roberts

---

## Abstract

We study F-rationality and F-regularity in diagonal subalgebras of multigraded rings, and use this to construct large families of rings that are F-rational but not F-regular. We also use diagonal subalgebras to construct rings with divisor class groups that are finitely generated but not discrete in the sense of Danilov. © 2008 Elsevier Inc. All rights reserved.

*Keywords:* Tight closure; Singularities; Diagonal subalgebras

---

## 1. Introduction

We study the properties of F-rationality and F-regularity in multigraded rings and their diagonal subalgebras. The main focus is on diagonal subalgebras of bigraded rings: these constitute an interesting class of rings since they arise naturally as homogeneous coordinate rings of blow-ups of projective varieties.

---

\* Corresponding author.

*E-mail addresses:* [kurano@math.meiji.ac.jp](mailto:kurano@math.meiji.ac.jp) (K. Kurano), [esato@math.kyushu-u.ac.jp](mailto:esato@math.kyushu-u.ac.jp) (E.-i. Sato), [singh@math.utah.edu](mailto:singh@math.utah.edu) (A.K. Singh), [watanabe@math.chs.nihon-u.ac.jp](mailto:watanabe@math.chs.nihon-u.ac.jp) (K.-i. Watanabe).

<sup>1</sup> Was supported by NSF grants DMS 0300600 and DMS 0600819.

Let  $X$  be a projective variety over a field  $K$ , with homogeneous coordinate ring  $A$ . Let  $\mathfrak{a} \subset A$  be a homogeneous ideal, and  $V \subset X$  the closed subvariety defined by  $\mathfrak{a}$ . For  $g$  an integer, we use  $\mathfrak{a}_g$  to denote the  $K$ -vector space consisting of homogeneous elements of  $\mathfrak{a}$  of degree  $g$ . If  $g \gg 0$ , then  $\mathfrak{a}_g$  defines a very ample complete linear system on the blow-up of  $X$  along  $V$ , and hence  $K[\mathfrak{a}_g]$  is a homogeneous coordinate ring for this blow-up. Since the ideals  $\mathfrak{a}^h$  define the same subvariety  $V$ , the rings  $K[(\mathfrak{a}^h)_g]$  are homogeneous coordinate ring for the blow-up provided  $g \gg h > 0$ .

Suppose that  $A$  is a standard  $\mathbb{N}$ -graded  $K$ -algebra, and consider the  $\mathbb{N}^2$ -grading on the Rees algebra  $A[at]$ , where  $\deg rt^j = (i, j)$  for  $r \in A_i$ . The connection with diagonal subalgebras stems from the fact that if  $\mathfrak{a}^h$  is generated by elements of degree less than or equal to  $g$ , then

$$K[(\mathfrak{a}^h)_g] \cong \bigoplus_{k \geq 0} A[at]_{(gk, hk)}.$$

Using  $\Delta = (g, h)\mathbb{Z}$  to denote the  $(g, h)$ -diagonal in  $\mathbb{Z}^2$ , the diagonal subalgebra  $A[at]_\Delta = \bigoplus_k A[at]_{(gk, hk)}$  is a homogeneous coordinate ring for the blow-up of  $\text{Proj } A$  along the subvariety defined by  $\mathfrak{a}$ , whenever  $g \gg h > 0$ .

The papers [GG,GGH,GGP,Tr] use diagonal subalgebras in studying blow-ups of projective space at finite sets of points. For  $A$  a polynomial ring and  $\mathfrak{a}$  a homogeneous ideal, the ring theoretic properties of  $K[\mathfrak{a}_g]$  are studied by Simis, Trung, and Valla in [STV] by realizing  $K[\mathfrak{a}_g]$  as a diagonal subalgebra of the Rees algebra  $A[at]$ . In particular, they determine when  $K[\mathfrak{a}_g]$  is Cohen–Macaulay for  $\mathfrak{a}$  a complete intersection ideal generated by forms of equal degree, and also for  $\mathfrak{a}$  the ideal of maximal minors of a generic matrix. Some of their results are extended by Conca, Herzog, Trung, and Valla as in the following theorem.

**Theorem 1.1.** (See [CHTV, Theorem 4.6].) *Let  $K[x_1, \dots, x_m]$  be a polynomial ring over a field, and let  $\mathfrak{a}$  be a complete intersection ideal minimally generated by forms of degrees  $d_1, \dots, d_r$ . Fix positive integers  $g$  and  $h$  with  $g/h > d = \max\{d_1, \dots, d_r\}$ .*

*Then  $K[(\mathfrak{a}^h)_g]$  is Cohen–Macaulay if and only if  $g > (h - 1)d - m + \sum_{j=1}^r d_j$ .*

When  $A$  is a polynomial ring and  $\mathfrak{a}$  an ideal for which  $A[at]$  is Cohen–Macaulay, Lavila-Vidal [Lv1, Theorem 4.5] proved that the diagonal subalgebras  $K[(\mathfrak{a}^h)_g]$  are Cohen–Macaulay for  $g \gg h \gg 0$ , thereby settling a conjecture from [CHTV]. In [CH] Cutkosky and Herzog obtain affirmative answers regarding the existence of a constant  $c$  such that  $K[(\mathfrak{a}^h)_g]$  is Cohen–Macaulay whenever  $g \geq ch$ . For more work on the Cohen–Macaulay and Gorenstein properties of diagonal subalgebras, see [HHR,Hy2,Lv2] and [LvZ].

As a motivating example for some of the results of this paper, consider a polynomial ring  $A = K[x_1, \dots, x_m]$  and an ideal  $\mathfrak{a} = (z_1, z_2)$  generated by relatively prime forms  $z_1$  and  $z_2$  of degree  $d$ . Setting  $\Delta = (d + 1, 1)\mathbb{Z}$ , the diagonal subalgebra  $A[at]_\Delta$  is a homogeneous coordinate ring for the blow-up of  $\text{Proj } A = \mathbb{P}^{m-1}$  along the subvariety defined by  $\mathfrak{a}$ . The Rees algebra  $A[at]$  has a presentation

$$\mathcal{R} = K[x_1, \dots, x_m, y_1, y_2]/(y_2z_1 - y_1z_2),$$

where  $\deg x_i = (1, 0)$  and  $\deg y_j = (d, 1)$ , and consequently  $\mathcal{R}_\Delta$  is the subalgebra of  $\mathcal{R}$  generated by the elements  $x_i y_j$ . When  $K$  has characteristic zero and  $z_1$  and  $z_2$  are general forms of degree  $d$ , the results of Section 3 imply that  $\mathcal{R}_\Delta$  has rational singularities if and only if  $d \leq m$ ,

and that it is of F-regular type if and only if  $d < m$ . As a consequence, we obtain large families of rings of the form  $\mathcal{R}_\Delta$ , standard graded over a field, which have rational singularities, but which are not of F-regular type.

It is worth pointing out that if  $\mathcal{R}$  is an  $\mathbb{N}^2$ -graded ring over an infinite field  $\mathcal{R}_{(0,0)} = K$ , and  $\Delta = (g, h)\mathbb{Z}$  for coprime positive integers  $g$  and  $h$ , then  $\mathcal{R}_\Delta$  is the ring of invariants of the torus  $K^*$  acting on  $\mathcal{R}$  via

$$\lambda : r \longmapsto \lambda^{hi-gj}r \quad \text{where } \lambda \in K^* \text{ and } r \in \mathcal{R}_{(i,j)}.$$

Consequently there exist torus actions on hypersurfaces for which the rings of invariants have rational singularities but are not of F-regular type.

In Section 4 we use diagonal subalgebras to construct standard graded normal rings  $R$ , with isolated singularities, for which  $H_m^2(R)_0 = 0$  and  $H_m^2(R)_1 \neq 0$ . If  $S$  is the localization of such a ring  $R$  at its homogeneous maximal ideal, then, by Danilov’s results, the divisor class group of  $S$  is a finitely generated abelian group, though  $S$  does not have a discrete divisor class group. Such rings  $R$  are also of interest in view of the results of [RSS], where it is proved that the image of  $H_m^2(R)_0$  in  $H_m^2(R^+)$  is annihilated by elements of  $R^+$  of arbitrarily small positive degree; here  $R^+$  denotes the absolute integral closure of  $R$ . A corresponding result for  $H_m^2(R)_1$  is not known at this point, and the rings constructed in Section 4 constitute interesting test cases.

Section 2 summarizes some notation and conventions for multigraded rings and modules. In Section 3 we carry out an analysis of diagonal subalgebras of bigraded hypersurfaces; this uses results on rational singularities and F-regular rings proved in Sections 5 and 6, respectively.

**2. Preliminaries**

In this section, we provide a brief treatment of multigraded rings and modules; see [GW1, GW2,HHR], and [HIO] for further details.

By an  $\mathbb{N}^r$ -graded ring we mean a ring

$$\mathcal{R} = \bigoplus_{\mathbf{n} \in \mathbb{N}^r} \mathcal{R}_{\mathbf{n}},$$

which is finitely generated over the subring  $\mathcal{R}_0$ . If  $(\mathcal{R}_0, \mathfrak{m})$  is a local ring, then  $\mathcal{R}$  has a unique homogeneous maximal ideal  $\mathfrak{M} = \mathfrak{m}\mathcal{R} + \mathcal{R}_+$ , where  $\mathcal{R}_+ = \bigoplus_{\mathbf{n} \neq 0} \mathcal{R}_{\mathbf{n}}$ .

For  $\mathbf{m} = (m_1, \dots, m_r)$  and  $\mathbf{n} = (n_1, \dots, n_r)$  in  $\mathbb{Z}^r$ , we say  $\mathbf{n} > \mathbf{m}$  (resp.  $\mathbf{n} \geq \mathbf{m}$ ) if  $n_i > m_i$  (resp.  $n_i \geq m_i$ ) for each  $i$ .

Let  $M$  be a  $\mathbb{Z}^r$ -graded  $\mathcal{R}$ -module. For  $\mathbf{m} \in \mathbb{Z}^r$ , we set

$$M_{\geq \mathbf{m}} = \bigoplus_{\mathbf{n} \geq \mathbf{m}} M_{\mathbf{n}},$$

which is a  $\mathbb{Z}^r$ -graded submodule of  $M$ . One writes  $M(\mathbf{m})$  for the  $\mathbb{Z}^r$ -graded  $\mathcal{R}$ -module with shifted grading  $[M(\mathbf{m})]_{\mathbf{n}} = M_{\mathbf{m}+\mathbf{n}}$  for each  $\mathbf{n} \in \mathbb{Z}^r$ .

Let  $M$  and  $N$  be  $\mathbb{Z}^r$ -graded  $\mathcal{R}$ -modules. Then  $\text{Hom}_{\mathcal{R}}(M, N)$  is the  $\mathbb{Z}^r$ -graded module with  $[\text{Hom}_{\mathcal{R}}(M, N)]_{\mathbf{n}}$  being the abelian group consisting of degree preserving  $\mathcal{R}$ -linear homomorphisms from  $M$  to  $N(\mathbf{n})$ .

The functor  $\underline{\text{Ext}}_{\mathcal{R}}^i(M, -)$  is the  $i$ th derived functor of  $\underline{\text{Hom}}_{\mathcal{R}}(M, -)$  in the category of  $\mathbb{Z}^r$ -graded  $\mathcal{R}$ -modules. When  $M$  is finitely generated,  $\underline{\text{Ext}}_{\mathcal{R}}^i(M, N)$  and  $\text{Ext}_{\mathcal{R}}^i(M, N)$  agree as underlying  $\mathcal{R}$ -modules. For a homogeneous ideal  $\mathfrak{a}$  of  $\mathcal{R}$ , the local cohomology modules of  $M$  with support in  $\mathfrak{a}$  are the  $\mathbb{Z}^r$ -graded modules

$$H_{\mathfrak{a}}^i(M) = \varinjlim_n \underline{\text{Ext}}_{\mathcal{R}}^i(\mathcal{R}/\mathfrak{a}^n, M).$$

Let  $\varphi : \mathbb{Z}^r \rightarrow \mathbb{Z}^s$  be a homomorphism of abelian groups satisfying  $\varphi(\mathbb{N}^r) \subseteq \mathbb{N}^s$ . We write  $\mathcal{R}^\varphi$  for the ring  $\mathcal{R}$  with the  $\mathbb{N}^s$ -grading where

$$[\mathcal{R}^\varphi]_n = \bigoplus_{\varphi(m)=n} \mathcal{R}_m.$$

If  $M$  is a  $\mathbb{Z}^r$ -graded  $\mathcal{R}$ -module, then  $M^\varphi$  is the  $\mathbb{Z}^s$ -graded  $\mathcal{R}^\varphi$ -module with

$$[M^\varphi]_n = \bigoplus_{\varphi(m)=n} M_m.$$

The change of grading functor  $(-)^{\varphi}$  is exact; by [HHR, Lemma 1.1] one has

$$H_{\mathfrak{M}}^i(M)^{\varphi} = H_{\mathfrak{M}^{\varphi}}^i(M^{\varphi}).$$

Consider the projections  $\varphi_i : \mathbb{Z}^r \rightarrow \mathbb{Z}$  with  $\varphi_i(m_1, \dots, m_r) = m_i$ , and set

$$a(\mathcal{R}^{\varphi_i}) = \max\{a \in \mathbb{Z} \mid [H_{\mathfrak{M}}^{\dim \mathcal{R}}(\mathcal{R})^{\varphi_i}]_a \neq 0\};$$

this is the  $a$ -invariant of the  $\mathbb{N}$ -graded ring  $\mathcal{R}^{\varphi_i}$  in the sense of Goto and Watanabe [GW1]. As in [HHR], the *multigraded  $\mathbf{a}$ -invariant* of  $\mathcal{R}$  is

$$\mathbf{a}(\mathcal{R}) = (a(\mathcal{R}^{\varphi_1}), \dots, a(\mathcal{R}^{\varphi_r})).$$

Let  $\mathcal{R}$  be a  $\mathbb{Z}^2$ -graded ring and let  $g, h$  be positive integers. The subgroup  $\Delta = (g, h)\mathbb{Z}$  is a *diagonal* in  $\mathbb{Z}^2$ , and the corresponding *diagonal subalgebra* of  $\mathcal{R}$  is

$$\mathcal{R}_{\Delta} = \bigoplus_{k \in \mathbb{Z}} \mathcal{R}_{(gk, hk)}.$$

Similarly, if  $M$  is a  $\mathbb{Z}^2$ -graded  $\mathcal{R}$ -module, we set

$$M_{\Delta} = \bigoplus_{k \in \mathbb{Z}} M_{(gk, hk)},$$

which is a  $\mathbb{Z}$ -graded module over the  $\mathbb{Z}$ -graded ring  $\mathcal{R}_{\Delta}$ .

**Lemma 2.1.** *Let  $A$  and  $B$  be  $\mathbb{N}$ -graded normal rings, finitely generated over a field  $A_0 = K = B_0$ . Set  $T = A \otimes_K B$ . Let  $g$  and  $h$  be positive integers and set  $\Delta = (g, h)\mathbb{Z}$ . Let  $\mathfrak{a}$ ,  $\mathfrak{b}$ , and  $\mathfrak{m}$  denote the homogeneous maximal ideals of  $A$ ,  $B$ , and  $T_\Delta$  respectively. Then, for each  $q \geq 0$  and  $i, j, k \in \mathbb{Z}$ , one has*

$$H_{\mathfrak{m}}^q(T(i, j)_\Delta)_k = (A_{i+gk} \otimes H_{\mathfrak{b}}^q(B)_{j+hk}) \oplus (H_{\mathfrak{a}}^q(A)_{i+gk} \otimes B_{j+hk}) \\ \oplus \bigoplus_{q_1+q_2=q+1} (H_{\mathfrak{a}}^{q_1}(A)_{i+gk} \otimes H_{\mathfrak{b}}^{q_2}(B)_{j+hk}).$$

**Proof.** Let  $A^{(g)}$  and  $B^{(h)}$  denote the respective Veronese subrings of  $A$  and  $B$ . Set

$$A^{(g,i)} = \bigoplus_{k \in \mathbb{Z}} A_{i+gk} \quad \text{and} \quad B^{(h,j)} = \bigoplus_{k \in \mathbb{Z}} B_{j+hk},$$

which are graded  $A^{(g)}$  and  $B^{(h)}$  modules respectively. Using  $\#$  for the Segre product,

$$T(i, j)_\Delta = \bigoplus_{k \in \mathbb{Z}} A_{i+gk} \otimes_K B_{j+hk} = A^{(g,i)} \# B^{(h,j)}.$$

The ideal  $A_+^{(g)}A$  is  $\mathfrak{a}$ -primary; likewise,  $B_+^{(h)}B$  is  $\mathfrak{b}$ -primary. The Künneth formula for local cohomology, [GW1, Theorem 4.1.5], now gives the desired result.  $\square$

**Notation 2.2.** *We use bold letters to denote lists of elements, e.g.,  $\mathbf{z} = z_1, \dots, z_s$  and  $\boldsymbol{\gamma} = \gamma_1, \dots, \gamma_s$ .*

### 3. Diagonal subalgebras of bigraded hypersurfaces

We prove the following theorem about diagonal subalgebras of  $\mathbb{N}^2$ -graded hypersurfaces. The proof uses results proved later in Sections 5 and 6.

**Theorem 3.1.** *Let  $K$  be a field, let  $m, n$  be integers with  $m, n \geq 2$ , and let*

$$\mathcal{R} = K[x_1, \dots, x_m, y_1, \dots, y_n]/(f)$$

*be a normal  $\mathbb{N}^2$ -graded hypersurface where  $\deg x_i = (1, 0)$ ,  $\deg y_j = (0, 1)$ , and  $\deg f = (d, e) > (0, 0)$ . For positive integers  $g$  and  $h$ , set  $\Delta = (g, h)\mathbb{Z}$ . Then:*

- (1) *The ring  $\mathcal{R}_\Delta$  is Cohen–Macaulay if and only if  $\lfloor (d - m)/g \rfloor < e/h$  and  $\lfloor (e - n)/h \rfloor < d/g$ . In particular, if  $d < m$  and  $e < n$ , then  $\mathcal{R}_\Delta$  is Cohen–Macaulay for each diagonal  $\Delta$ .*
- (2) *The graded canonical module of  $\mathcal{R}_\Delta$  is  $\mathcal{R}(d - m, e - n)_\Delta$ . Hence  $\mathcal{R}_\Delta$  is Gorenstein if and only if  $(d - m)/g = (e - n)/h$ , and this is an integer.*

*If  $K$  has characteristic zero, and  $f$  is a generic polynomial of degree  $(d, e)$ , then:*

- (3) *The ring  $\mathcal{R}_\Delta$  has rational singularities if and only if it is Cohen–Macaulay and  $d < m$  or  $e < n$ .*
- (4) *The ring  $\mathcal{R}_\Delta$  is of  $F$ -regular type if and only if  $d < m$  and  $e < n$ .*




Fig. 1. Properties of  $\mathcal{R}_\Delta$  for  $\Delta = (1, 1)\mathbb{Z}$ .

For  $m, n \geq 3$  and  $\Delta = (1, 1)\mathbb{Z}$ , the properties of  $\mathcal{R}_\Delta$ , as determined by  $m, n, d, e$ , are summarized in Fig. 1.

**Remark 3.2.** Let  $m, n \geq 2$ . A generic hypersurface of degree  $(d, e) > (0, 0)$  in  $m, n$  variables is normal precisely when

$$m > \min(2, d) \quad \text{and} \quad n > \min(2, e).$$

Suppose that  $m = 2 = n$ , and that  $f$  is nonzero. Then  $\dim \mathcal{R}_\Delta = 2$ ; since  $\mathcal{R}_\Delta$  is generated over a field by elements of equal degree,  $\mathcal{R}_\Delta$  is of F-regular type if and only if it has rational singularities; see [Wa2]. This is the case precisely if

$$\begin{aligned} d = 1, \quad e \leq h + 1, \quad \text{or} \\ e = 1, \quad d \leq g + 1. \end{aligned}$$

Following a suggestion of Hara, the case  $n = 2$  and  $e = 1$  was used in [Si, Example 7.3] to construct examples of standard graded rings with rational singularities which are not of F-regular type.

**Proof of Theorem 3.1.** Set  $A = K[x]$ ,  $B = K[y]$ , and  $T = A \otimes_K B$ . By Lemma 2.1,  $H_m^q(T_\Delta) = 0$  for  $q \neq m + n - 1$ . The local cohomology exact sequence induced by

$$0 \longrightarrow T(-d, -e)_\Delta \xrightarrow{f} T_\Delta \longrightarrow \mathcal{R}_\Delta \longrightarrow 0$$

therefore gives  $H_m^{q-1}(\mathcal{R}_\Delta) = H_m^q(T(-d, -e)_\Delta)$  for  $q \leq m + n - 2$ , and also shows that  $H_m^{m+n-2}(\mathcal{R}_\Delta)$  and  $H_m^{m+n-1}(\mathcal{R}_\Delta)$  are, respectively, the kernel and cokernel of

$$\begin{array}{ccc} H_m^{m+n-1}(T(-d, -e)_\Delta) & \xrightarrow{f} & H_m^{m+n-1}(T_\Delta) \\ \parallel & & \parallel \\ [H_a^m(A(-d)) \otimes H_b^n(B(-e))]_\Delta & \xrightarrow{f} & [H_a^m(A) \otimes H_b^n(B)]_\Delta. \end{array}$$

The horizontal map above is surjective since its graded dual

$$\begin{array}{ccc} [A(d-m) \otimes B(e-n)]_\Delta & \xleftarrow{f} & [A(-m) \otimes B(-n)]_\Delta \\ \parallel & & \parallel \\ T(d-m, e-n)_\Delta & \xleftarrow{f} & T(-m, -n)_\Delta \end{array}$$

is injective. In particular,  $\dim \mathcal{R}_\Delta = m + n - 2$ .

It follows from the above discussion that  $\mathcal{R}_\Delta$  is Cohen–Macaulay if and only if  $H_m^q(T(-d, -e)_\Delta) = 0$  for each  $q \leq m + n - 2$ . By Lemma 2.1, this is the case if and only if, for each integer  $k$ , one has

$$A_{-d+gk} \otimes H_b^n(B)_{-e+hk} = 0 = H_a^m(A)_{-d+gk} \otimes B_{-e+hk}.$$

Hence  $\mathcal{R}_\Delta$  is Cohen–Macaulay if and only if there is no integer  $k$  satisfying

$$d/g \leq k \leq (e-n)/h \quad \text{or} \quad e/h \leq k \leq (d-m)/g,$$

which completes the proof of (1).

For (2), note that the graded canonical module of  $\mathcal{R}_\Delta$  is the graded dual of  $H_m^{m+n-2}(\mathcal{R}_\Delta)$ , and hence that it equals

$$\text{coker}(T(-m, -n)_\Delta \xrightarrow{f} T(d-m, e-n)_\Delta) = \mathcal{R}(d-m, e-n)_\Delta.$$

This module is principal if and only if  $\mathcal{R}(d-m, e-n)_\Delta = \mathcal{R}_\Delta(a)$  for some integer  $a$ , i.e.,  $d-m = ga$  and  $e-n = ha$ .

When  $f$  is a general polynomial of degree  $(d, e)$ , the ring  $\mathcal{R}_\Delta$  has an isolated singularity. Also,  $\mathcal{R}_\Delta$  is normal since it is a direct summand of the normal ring  $\mathcal{R}$ . By Theorem 5.1,  $\mathcal{R}_\Delta$  has rational singularities precisely if it is Cohen–Macaulay and  $a(\mathcal{R}_\Delta) < 0$ ; this proves (3).

It remains to prove (4). If  $d < m$  and  $e < n$ , then Theorem 5.2 implies that  $\mathcal{R}$  has rational singularities. By Theorem 6.2, it follows that for almost all primes  $p$ , the characteristic  $p$  models  $\mathcal{R}_p$  of  $\mathcal{R}$  are F-rational hypersurfaces which, therefore, are F-regular. Alternatively,  $\mathcal{R}_p$  is a generic hypersurface of degree  $(d, e) < (m, n)$ , so Theorem 6.5 implies that  $\mathcal{R}_p$  is F-regular. Since  $(\mathcal{R}_p)_\Delta$  is a direct summand of  $\mathcal{R}_p$ , it follows that  $(\mathcal{R}_p)_\Delta$  is F-regular. The rings  $(\mathcal{R}_p)_\Delta$  are characteristic  $p$  models of  $\mathcal{R}_\Delta$ , so we conclude that  $\mathcal{R}_\Delta$  is of F-regular type.

Suppose  $\mathcal{R}_\Delta$  has F-regular type, and let  $(\mathcal{R}_p)_\Delta$  be a characteristic  $p$  model which is F-regular. Fix an integer  $k > d/g$ . Then Proposition 6.3 implies that there exists an integer  $q = p^e$  such that

$$\text{rank}_K ((\mathcal{R}_p)_\Delta)_k \leq \text{rank}_K [H_m^{m+n-2}(\omega^{(q)})]_k,$$

where  $\omega$  is the graded canonical module of  $(\mathcal{R}_p)_\Delta$ . Using (2), we see that

$$H_m^{m+n-2}(\omega^{(q)}) = H_m^{m+n-2}(\mathcal{R}_p(qd - qm, qe - qn)_\Delta).$$

Let  $T_p$  be a characteristic  $p$  model for  $T$  such that  $T_p/fT_p = \mathcal{R}_p$ . Multiplication by  $f$  on  $T_p$  induces a local cohomology exact sequence

$$\begin{aligned} \dots \longrightarrow H_{m_p}^{m+n-2}(T_p(qd - qm, qe - qn)_\Delta) &\longrightarrow H_{m_p}^{m+n-2}(\mathcal{R}_p(qd - qm, qe - qn)_\Delta) \\ &\longrightarrow H_{m_p}^{m+n-1}(T_p(qd - qm - d, qe - qn - e)_\Delta) \longrightarrow \dots \end{aligned}$$

Since  $H_{m_p}^{m+n-2}(T_p(qd - qm, qe - qn)_\Delta)$  vanishes by Lemma 2.1, we conclude that

$$\begin{aligned} \text{rank}_K ((\mathcal{R}_p)_\Delta)_k &\leq \text{rank}_K [H_{m_p}^{m+n-1}(T_p(qd - qm - d, qe - qn - e)_\Delta)]_k \\ &= \text{rank}_K H_{a_p}^m(A_p)_{qd - qm - d + gk} \otimes H_{b_q}^n(B_p)_{qe - qn - e + hk}. \end{aligned}$$

Hence  $qd - qm - d + gk < 0$ ; as  $d - gk < 0$ , we conclude  $d < m$ . Similarly,  $e < n$ .  $\square$

We conclude this section with an example where a local cohomology module of a standard graded ring is not rigid in the sense that  $H_m^2(R)_0 = 0$  while  $H_m^2(R)_1 \neq 0$ . Further such examples are constructed in Section 4.

**Proposition 3.3.** *Let  $K$  be a field and let*

$$\mathcal{R} = K[x_1, x_2, x_3, y_1, y_2]/(f)$$

where  $\deg x_i = (1, 0)$ ,  $\deg y_j = (0, 1)$ , and  $\deg f = (d, e)$  for  $d \geq 4$  and  $e \geq 1$ . Let  $g$  and  $h$  be positive integers such that  $g \leq d - 3$  and  $h \geq e$ , and set  $\Delta = (g, h)\mathbb{Z}$ . Then  $H_m^2(\mathcal{R}_\Delta)_0 = 0$  and  $H_m^2(\mathcal{R}_\Delta)_1 \neq 0$ .

**Proof.** Using the resolution of  $\mathcal{R}$  over the polynomial ring  $T$  as in the proof of Theorem 3.1, we have an exact sequence

$$H_m^2(T_\Delta) \longrightarrow H_m^2(\mathcal{R}_\Delta) \longrightarrow H_m^3(T(-d, -e)_\Delta) \longrightarrow H_m^3(T_\Delta).$$

Lemma 2.1 implies that  $H_m^2(T_\Delta) = 0 = H_m^3(T_\Delta)$ . Hence, again by Lemma 2.1,

$$H_m^2(\mathcal{R}_\Delta)_0 = H^3(A)_{-d} \otimes B_{-e} = 0 \quad \text{and} \quad H_m^2(\mathcal{R}_\Delta)_1 = H^3(A)_{g-d} \otimes B_{h-e} \neq 0. \quad \square$$



### 4. Non-rigid local cohomology modules

We construct examples of standard graded normal rings  $R$  over  $\mathbb{C}$ , with only isolated singularities, for which  $H_m^2(R)_0 = 0$  and  $H_m^2(R)_1 \neq 0$ . Let  $S$  be the localization of such a ring  $R$  at its homogeneous maximal ideal. By results of Danilov [Da1, Da2], Theorem 4.1 below, it follows that the divisor class group of  $S$  is finitely generated, though  $S$  does not have a discrete divisor class group, i.e., the natural map  $\text{Cl}(S) \rightarrow \text{Cl}(S[[t]])$  is not bijective. Here, remember that if  $A$  is a Noetherian normal domain, then so is  $A[[t]]$ .

**Theorem 4.1.** *Let  $R$  be a standard graded normal ring, which is finitely generated as an algebra over  $R_0 = \mathbb{C}$ . Assume, moreover, that  $X = \text{Proj } R$  is smooth. Set  $(S, \mathfrak{m})$  to be the local ring of  $R$  at its homogeneous maximal ideal, and  $\widehat{S}$  to be the  $\mathfrak{m}$ -adic completion of  $S$ . Then*

- (1) *the group  $\text{Cl}(S)$  is finitely generated if and only if  $H^1(X, \mathcal{O}_X) = 0$ ;*
- (2) *the map  $\text{Cl}(S) \rightarrow \text{Cl}(\widehat{S})$  is bijective if and only if  $H^1(X, \mathcal{O}_X(i)) = 0$  for each integer  $i \geq 1$ ;*  
*and*
- (3) *the map  $\text{Cl}(S) \rightarrow \text{Cl}(S[[t]])$  is bijective if and only if  $H^1(X, \mathcal{O}_X(i)) = 0$  for each integer  $i \geq 0$ .*

The essential point in our construction is in the following theorem.

**Theorem 4.2.** *Let  $A$  be a Cohen–Macaulay ring of dimension  $d \geq 2$ , which is a standard graded algebra over a field  $K$ . For  $s \geq 2$ , let  $z_1, \dots, z_s$  be a regular sequence in  $A$ , consisting of homogeneous elements of equal degree, say  $k$ . Consider the Rees ring  $\mathcal{R} = A[z_1t, \dots, z_st]$  with the  $\mathbb{Z}^2$ -grading where  $\deg x = (n, 0)$  for  $x \in A_n$ , and  $\deg z_it = (0, 1)$ .*

*Let  $\Delta = (g, h)\mathbb{Z}$  where  $g, h$  are positive integers, and let  $\mathfrak{m}$  denote the homogeneous maximal ideal of  $\mathcal{R}_\Delta$ . Then:*

- (1)  *$H_m^q(\mathcal{R}_\Delta) = 0$  if  $q \neq d - s + 1, d$ ; and*
- (2)  *$H_m^{d-s+1}(\mathcal{R}_\Delta)_i \neq 0$  if and only if  $1 \leq i \leq (a + ks - k)/g$ , where  $a$  is the  $a$ -invariant of  $A$ .*

*In particular,  $\mathcal{R}_\Delta$  is Cohen–Macaulay if and only if  $g > a + ks - k$ .*

**Example 4.3.** For  $d \geq 3$ , let  $A = \mathbb{C}[x_0, \dots, x_d]/(f)$  be a standard graded hypersurface such that  $\text{Proj } A$  is smooth over  $\mathbb{C}$ . Take general  $k$ -forms  $z_1, \dots, z_{d-1} \in A$ , and consider the Rees ring  $\mathcal{R} = A[z_1t, \dots, z_{d-1}t]$ . Since  $(z) \subset A$  is a radical ideal,

$$\text{gr}((z), A) \cong A/(z)[y_1, \dots, y_{d-1}]$$

is a reduced ring, and therefore  $\mathcal{R} = A[z_1t, \dots, z_{d-1}t]$  is integrally closed in  $A[t]$ . Since  $A$  is normal, so is  $\mathcal{R}$ . Note that  $\text{Proj } \mathcal{R}_\Delta$  is the blow-up of  $\text{Proj } A$  at the subvariety defined by  $(z)$ , i.e., at  $k^{d-1}(\deg f)$  points. It follows that  $\text{Proj } \mathcal{R}_\Delta$  is smooth over  $\mathbb{C}$ . Hence  $\mathcal{R}_\Delta$  is a standard graded  $\mathbb{C}$ -algebra, which is normal and has an isolated singularity.

If  $\Delta = (g, h)\mathbb{Z}$  is a diagonal with  $1 \leq g \leq \deg f + k(d - 2) - (d + 1)$  and  $h \geq 1$ , then Theorem 4.2 implies that

$$H_m^2(\mathcal{R}_\Delta)_0 = 0 \quad \text{and} \quad H_m^2(\mathcal{R}_\Delta)_1 \neq 0.$$

The rest of this section is devoted to proving Theorem 4.2. We may assume that the base field  $K$  is infinite. Then one can find linear forms  $x_1, \dots, x_{d-s}$  in  $A$  such that  $x_1, \dots, x_{d-s}, z_1, \dots, z_s$  is a maximal  $A$ -regular sequence.

We will use the following lemma; the notation is as in Theorem 4.2.

**Lemma 4.4.** *Let  $\mathfrak{a}$  be the homogeneous maximal ideal of  $A$ . Set  $I = (z_1, \dots, z_s)A$ . Let  $r$  be a positive integer.*

- (1)  $H_{\mathfrak{a}}^q(I^r) = 0$  if  $q \neq d - s + 1, d$ .
- (2) Assume  $d > s$ . Then,  $H_{\mathfrak{a}}^{d-s+1}(I^r)_i \neq 0$  if and only if  $i \leq a + ks + rk - k$ .
- (3) Assume  $d = s$ . Then,  $H_{\mathfrak{a}}^{d-s+1}(I^r)_i \neq 0$  if and only if  $0 \leq i \leq a + ks + rk - k$ .

**Proof.** Recall that  $A$  and  $A/I^r$  are Cohen–Macaulay rings of dimension  $d$  and  $d - s$ , respectively. By the exact sequence

$$0 \longrightarrow I^r \longrightarrow A \longrightarrow A/I^r \longrightarrow 0$$

we obtain

$$H_{\mathfrak{a}}^q(I^r) = \begin{cases} H_{\mathfrak{a}}^d(A) & \text{if } q = d, \\ H_{\mathfrak{a}}^{d-s}(A/I^r) & \text{if } q = d - s + 1, \\ 0 & \text{if } q \neq d - s + 1, d, \end{cases}$$

which proves (1).

Next we prove (2) and (3). Since  $A/I^r$  is a standard graded Cohen–Macaulay ring of dimension  $d - s$ , it is enough to show that the  $a$ -invariant of this ring equals  $a + ks + rk - k$ . This is straightforward if  $r = 1$ , and we proceed by induction. Consider the exact sequence

$$0 \longrightarrow I^r/I^{r+1} \longrightarrow A/I^{r+1} \longrightarrow A/I^r \longrightarrow 0.$$

Since  $z_1, \dots, z_s$  is a regular sequence of  $k$ -forms,  $I^r/I^{r+1}$  is isomorphic to

$$((A/I)(-rk))^{\binom{s-1+r}{r}}.$$

Thus, we have the following exact sequence:

$$0 \longrightarrow H_{\mathfrak{a}}^{d-s}((A/I)(-rk))^{\binom{s-1+r}{r}} \longrightarrow H_{\mathfrak{a}}^{d-s}(A/I^{r+1}) \longrightarrow H_{\mathfrak{a}}^{d-s}(A/I^r) \longrightarrow 0.$$

The  $a$ -invariant of  $(A/I)(-rk)$  equals  $a + ks + rk$ , and that of  $A/I^r$  is  $a + ks + rk - k$  by the inductive hypothesis. Thus,  $A/I^{r+1}$  has  $a$ -invariant  $a + ks + rk$ .  $\square$

**Proof of Theorem 4.2.** Let  $B = K[y_1, \dots, y_s]$  be a polynomial ring, and set

$$T = A \otimes_K B = A[y_1, \dots, y_s].$$

Consider the  $\mathbb{Z}^2$ -grading on  $T$  where  $\deg x = (n, 0)$  for  $x \in A_n$ , and  $\deg y_i = (0, 1)$  for each  $i$ . One has a surjective homomorphism of graded rings

$$T \longrightarrow \mathcal{R} = A[z_1t, \dots, z_s t] \quad \text{where } y_i \longmapsto z_i t,$$

and this induces an isomorphism

$$\mathcal{R} \cong T/I_2 \begin{pmatrix} z_1 & \cdots & z_s \\ y_1 & \cdots & y_s \end{pmatrix}.$$

The minimal free resolution of  $\mathcal{R}$  over  $T$  is given by the Eagon–Northcott complex

$$0 \longrightarrow F^{-(s-1)} \longrightarrow F^{-(s-2)} \longrightarrow \dots \longrightarrow F^0 \longrightarrow 0,$$

where  $F^0 = T(0, 0)$ , and  $F^{-i}$  for  $1 \leq i \leq s - 1$  is the direct sum of  $\binom{s}{i+1}$  copies of

$$T(-k, -i) \oplus T(-2k, -(i - 1)) \oplus \dots \oplus T(-ik, -1).$$

Let  $\mathfrak{n}$  be the homogeneous maximal ideal of  $T_\Delta$ . One has the spectral sequence:

$$E_2^{p,q} = H^p(H_{\mathfrak{n}}^q(F_\Delta^\bullet)) \implies H_{\mathfrak{m}}^{p+q}(\mathcal{R}_\Delta).$$

Let  $G$  be the set of  $(n, m)$  such that  $T(n, m)$  appears in the Eagon–Northcott complex above, i.e., the elements of  $G$  are

$$\begin{aligned} &(0, 0), \\ &(-k, -1), \\ &(-k, -2), (-2k, -1), \\ &(-k, -3), (-2k, -2), (-3k, -1), \\ &\vdots \\ &(-k, -(s - 1)), \dots, (-(s - 1)k, -1). \end{aligned}$$

Let  $\mathfrak{a}$  and  $\mathfrak{b}$  be the homogeneous maximal ideal of  $A$  and  $B$ , respectively. For integers  $n$  and  $m$ , the Künneth formula gives

$$\begin{aligned} &H_{\mathfrak{n}}^q(T(n, m)) \\ &= H_{\mathfrak{n}}^q(A(n) \otimes_K B(m)) \\ &= (H_{\mathfrak{a}}^q(A(n)) \otimes B(m)) \oplus (A(n) \otimes H_{\mathfrak{b}}^q(B(m))) \oplus \bigoplus_{i+j=q+1} H_{\mathfrak{a}}^i(A(n)) \otimes H_{\mathfrak{b}}^j(B(m)) \\ &= H_{\mathfrak{a}}^q(T(n, m)) \oplus H_{\mathfrak{b}}^q(T(n, m)) \oplus \bigoplus_{i+j=q+1} H_{\mathfrak{a}}^i(A(n)) \otimes_K H_{\mathfrak{b}}^j(B(m)). \end{aligned}$$

As  $A$  and  $B$  are Cohen–Macaulay of dimension  $d$  and  $s$  respectively, it follows that

$$H_{\mathfrak{n}}^q(F^\bullet) = 0 \quad \text{if } q \neq s, d, d + s - 1.$$

In the case where  $d > s$ , one has

$$H_n^s(F^\bullet) = H_b^s(F^\bullet) \quad \text{and} \quad H_n^d(F^\bullet) = H_a^d(F^\bullet),$$

and if  $d = s$ , then

$$H_n^d(F^\bullet) = H_a^d(F^\bullet) \oplus H_b^s(F^\bullet).$$

We claim  $H_b^s(F^\bullet)_\Delta = 0$ . If not, there exists  $(n, m) \in G$  and  $\ell \in \mathbb{Z}$  such that

$$H_b^s(T(n, m))_{(g\ell, h\ell)} \neq 0.$$

This implies that

$$H_b^s(T(n, m))_{(g\ell, h\ell)} = A(n)_{g\ell} \otimes_K H_b^s(B(m))_{h\ell} = A_{n+g\ell} \otimes_K H_b^s(B)_{m+h\ell}$$

is nonzero, so

$$n + g\ell \geq 0 \quad \text{and} \quad m + h\ell \leq -s,$$

and hence

$$-\frac{n}{g} \leq \ell \leq -\frac{s+m}{h}.$$

But  $(n, m) \in G$ , so  $n \leq 0$  and  $m \geq -(s - 1)$ , implying that

$$0 \leq \ell \leq -\frac{1}{h},$$

which is not possible. This proves that  $H_b^s(F^\bullet)_\Delta = 0$ . Thus, we have

$$H_n^q(F^\bullet)_\Delta = \begin{cases} 0 & \text{if } q \neq d, d + s - 1, \\ H_a^d(F^\bullet)_\Delta & \text{if } q = d. \end{cases}$$

It follows that

$$E_2^{p,q} = H^p(H_n^q(F^\bullet)_\Delta) = E_\infty^{p,q}$$

for each  $p$  and  $q$ . Therefore,

$$H_m^i(\mathcal{R}_\Delta) = E_2^{i-d,d} = H^{i-d}(H_n^d(F^\bullet)_\Delta) = H^{i-d}(H_a^d(F^\bullet)_\Delta) = H_a^i(\mathcal{R})_\Delta$$

for  $d - s + 1 \leq i \leq d - 1$ , and

$$H_m^i(\mathcal{R}_\Delta) = 0 \quad \text{for } i < d - s + 1.$$

We next study  $H_a^i(\mathcal{R})$ . Since

$$\mathcal{R} = A \oplus I(k) \oplus I^2(2k) \oplus \cdots \oplus I^r(rk) \oplus \cdots,$$

we have

$$H_a^i(\mathcal{R}) = H_a^i(A) \oplus H_a^i(I)(k) \oplus H_a^i(I^2)(2k) \oplus \cdots \oplus H_a^i(I^r)(rk) \oplus \cdots.$$

Theorem 4.2 (1) now follow using Lemma 4.4 (1).

Assume that  $d > s$ . Then, by Lemma 4.4 (2),  $H_a^{d-s+1}(I^r(rk))_i \neq 0$  if and only if  $i \leq a + ks - k$ .

Assume that  $d = s$ . Then, by Lemma 4.4 (3),  $H_a^{d-s+1}(I^r(rk))_i \neq 0$  if and only if  $-rk \leq i \leq a + ks - k$ .

In each case,  $H_a^{d-s+1}(\mathcal{R})_{(gi,hi)} \neq 0$  if and only if

$$1 \leq i \leq \frac{a + ks - k}{g}. \quad \square$$

### 5. Rational singularities

Let  $R$  be a normal domain, essentially of finite type over a field of characteristic zero, and consider a *desingularization*  $f : Z \rightarrow \text{Spec } R$ , i.e., a proper birational morphism with  $Z$  a non-singular variety. One says  $R$  has *rational singularities* if  $R^i f_* \mathcal{O}_Z = 0$  for each  $i \geq 1$ ; this does not depend on the choice of the desingularization  $f$ . For  $\mathbb{N}$ -graded rings, one has the following criterion due to Flenner [Fl] and Watanabe [Wa1].

**Theorem 5.1.** *Let  $R$  be a normal  $\mathbb{N}$ -graded ring which is finitely generated over a field  $R_0$  of characteristic zero. Then  $R$  has rational singularities if and only if it is Cohen–Macaulay,  $a(R) < 0$ , and the localization  $R_{\mathfrak{p}}$  has rational singularities for each  $\mathfrak{p} \in \text{Spec } R \setminus \{R_+\}$ .*

When  $R$  has an isolated singularity, the above theorem gives an effective criterion for determining if  $R$  has rational singularities. However, a multigraded hypersurface typically does not have an isolated singularity, and the following variation turns out to be useful.

**Theorem 5.2.** *Let  $R$  be a normal  $\mathbb{N}^r$ -graded ring such that  $R_0$  is a local ring essentially of finite type over a field of characteristic zero, and  $R$  is generated over  $R_0$  by elements*

$$x_{11}, x_{12}, \dots, x_{1t_1}, \quad x_{21}, x_{22}, \dots, x_{2t_2}, \quad \dots, \quad x_{r1}, x_{r2}, \dots, x_{rt_r},$$

where  $\deg x_{ij}$  is a positive integer multiple of the  $i$ th unit vector  $e_i \in \mathbb{N}^r$ . Then  $R$  has rational singularities if and only if

- (1)  $R$  is Cohen–Macaulay,
- (2)  $R_{\mathfrak{p}}$  has rational singularities for each  $\mathfrak{p}$  belonging to

$$\text{Spec } R \setminus (V(x_{11}, x_{12}, \dots, x_{1t_1}) \cup \cdots \cup V(x_{r1}, x_{r2}, \dots, x_{rt_r})), \quad \text{and}$$

(3)  $a(R) < \mathbf{0}$ , i.e.,  $a(R^{\varphi_i}) < 0$  for each coordinate projection  $\varphi_i : \mathbb{N}^r \rightarrow \mathbb{N}$ .

Before proceeding with the proof, we record some preliminary results.

**Remark 5.3.** Let  $R$  be an  $\mathbb{N}$ -graded ring. We use  $R^{\natural}$  to denote the Rees algebra with respect to the filtration  $F_n = R_{\geq n}$ , i.e.,

$$R^{\natural} = F_0 \oplus F_1 T \oplus F_2 T^2 \oplus \dots$$

When considering  $\text{Proj } R^{\natural}$ , we use the  $\mathbb{N}$ -grading on  $R^{\natural}$  where  $[R^{\natural}]_n = F_n T^n$ . The inclusion  $R = [R^{\natural}]_0 \hookrightarrow R^{\natural}$  gives a map

$$\text{Proj } R^{\natural} \xrightarrow{f} \text{Spec } R.$$

Also, the inclusions  $R_n \hookrightarrow F_n$  give rise to an injective homomorphism of graded rings  $R \hookrightarrow R^{\natural}$ , which induces a surjection

$$\text{Proj } R^{\natural} \xrightarrow{\pi} \text{Proj } R.$$

**Lemma 5.4.** Let  $R$  be an  $\mathbb{N}$ -graded ring which is finitely generated over  $R_0$ , and assume that  $R_0$  is essentially of finite type over a field of characteristic zero.

If  $R_{\mathfrak{p}}$  has rational singularities for all primes  $\mathfrak{p} \in \text{Spec } R \setminus V(R_+)$ , then  $\text{Proj } R^{\natural}$  has rational singularities.

**Proof.** Note that  $\text{Proj } R^{\natural}$  is covered by affine open sets  $D_+(rT^n)$  for integers  $n \geq 1$  and homogeneous elements  $r \in R_{\geq n}$ . Consequently, it suffices to check that  $[R_{rT^n}^{\natural}]_0$  has rational singularities. Next, note that

$$[R_{rT^n}^{\natural}]_0 = R + \frac{1}{r}[R]_{\geq n} + \frac{1}{r^2}[R]_{\geq 2n} + \dots$$

In the case  $\text{deg } r > n$ , the ring above is simply  $R_r$ , which has rational singularities by the hypothesis of the lemma. If  $\text{deg } r = n$ , then

$$[R_{rT^n}^{\natural}]_0 = [R_r]_{\geq 0}.$$

The  $\mathbb{Z}$ -graded ring  $R_r$  has rational singularities and so, by [Wa1, Lemma 2.5], the ring  $[R_r]_{\geq 0}$  has rational singularities as well.  $\square$

**Lemma 5.5.** (See [Hy2, Lemma 2.3].) Let  $R$  be an  $\mathbb{N}$ -graded ring which is finitely generated over a local ring  $(R_0, \mathfrak{m})$ . Suppose  $[H_{\mathfrak{m}+R_+}^i(R)]_{\geq 0} = 0$  for all  $i \geq 0$ . Then, for all ideals  $\mathfrak{a}$  of  $R_0$ , one has

$$[H_{\mathfrak{a}+R_+}^i(R)]_{\geq 0} = 0 \quad \text{for all } i \geq 0.$$

We are now in a position to prove the following theorem, which is a variation of [Fl, Satz 3.1], [Wa1, Theorem 2.2], and [Hy1, Theorem 1.5].

**Theorem 5.6.** *Let  $R$  be an  $\mathbb{N}$ -graded normal ring which is finitely generated over  $R_0$ , and assume that  $R_0$  is a local ring essentially of finite type over a field of characteristic zero. Then  $R$  has rational singularities if and only if*

- (1)  $R$  is Cohen–Macaulay,
- (2)  $R_{\mathfrak{p}}$  has rational singularities for all  $\mathfrak{p} \in \text{Spec } R \setminus V(R_+)$ , and
- (3)  $a(R) < 0$ .

**Proof.** It is straightforward to see that conditions (1)–(3) hold when  $R$  has rational singularities, and we focus on the converse. Consider the morphism

$$Y = \text{Proj } R^{\natural} \xrightarrow{f} \text{Spec } R$$

as in Remark 5.3. Let  $g : Z \rightarrow Y$  be a desingularization of  $Y$ ; the composition

$$Z \xrightarrow{g} Y \xrightarrow{f} \text{Spec } R$$

is then a desingularization of  $\text{Spec } R$ . Note that  $Y = \text{Proj } R^{\natural}$  has rational singularities by Lemma 5.4, so

$$g_*\mathcal{O}_Z = \mathcal{O}_Y \quad \text{and} \quad R^q g_*\mathcal{O}_Z = 0 \quad \text{for all } q \geq 1.$$

Consequently the Leray spectral sequence

$$E_2^{p,q} = H^p(Y, R^q g_*\mathcal{O}_Z) \implies H^{p+q}(Z, \mathcal{O}_Z)$$

degenerates, and we get  $H^p(Z, \mathcal{O}_Z) = H^p(Y, \mathcal{O}_Y)$  for all  $p \geq 1$ . Since  $\text{Spec } R$  is affine, we also have  $R^p(g \circ f)_*\mathcal{O}_Z = H^p(Z, \mathcal{O}_Z)$ . To prove that  $R$  has rational singularities, it now suffices to show that  $H^p(Y, \mathcal{O}_Y) = 0$  for all  $p \geq 1$ . Consider the map  $\pi : Y \rightarrow X = \text{Proj } R$ . We have

$$H^p(Y, \mathcal{O}_Y) = H^p(X, \pi_*\mathcal{O}_X) = \bigoplus_{n \geq 0} H^p(X, \mathcal{O}_X(n)) = [H_{R_+}^{p+1}(R)]_{\geq 0}.$$

By condition (1), we have  $[H_{\mathfrak{m}+R_+}^p(R)]_{\geq 0} = 0$  for all  $p \geq 0$ , and so Lemma 5.5 implies that  $[H_{R_+}^p(R)]_{\geq 0} = 0$  for all  $p \geq 0$  as desired.  $\square$

**Proof of theorem 5.2.** If  $R$  has rational singularities, it is easily seen that conditions (1)–(3) must hold. For the converse, we proceed by induction on  $r$ . The case  $r = 1$  is Theorem 5.6 established above, so assume  $r \geq 2$ . It suffices to show that  $R_{\mathfrak{M}}$  has rational singularities where  $\mathfrak{M}$  is the homogeneous maximal ideal of  $R$ . Set

$$\mathfrak{m} = \mathfrak{M} \cap [R^{\varphi_r}]_0,$$

and consider the  $\mathbb{N}$ -graded ring  $S$  obtained by inverting the multiplicative set  $[R^{\varphi_r}]_0 \setminus \mathfrak{m}$  in  $R^{\varphi_r}$ . Since  $R_{\mathfrak{M}}$  is a localization of  $S$ , it suffices to show that  $S$  has rational singularities. Note that

$a(S) = a(R^{\varphi_r})$ , which is a negative integer by (1). Using Theorem 5.6, it is therefore enough to show that  $R_{\mathfrak{P}}$  has rational singularities for all  $\mathfrak{P} \in \text{Spec } R \setminus V(x_{r_1}, x_{r_2}, \dots, x_{r_t_r})$ . Fix such a prime  $\mathfrak{P}$ , and let

$$\psi : \mathbb{Z}^r \longrightarrow \mathbb{Z}^{r-1}$$

be the projection to the first  $r - 1$  coordinates. Note that  $R^\psi$  is the ring  $R$  regraded such that  $\text{deg } x_{r_j} = 0$ , and the degrees of  $x_{ij}$  for  $i < r$  are unchanged. Set

$$\mathfrak{p} = \mathfrak{P} \cap [R^\psi]_{\mathbf{0}},$$

and let  $T$  be the ring obtained by inverting the multiplicative set  $[R^\psi]_{\mathbf{0}} \setminus \mathfrak{p}$  in  $R^\psi$ . It suffices to show that  $T$  has rational singularities. Note that  $T$  is an  $\mathbb{N}^{r-1}$ -graded ring defined over a local ring  $(T_{\mathbf{0}}, \mathfrak{p})$ , and that it has homogeneous maximal ideal  $\mathfrak{p} + \mathfrak{b}T$  where

$$\mathfrak{b} = (R^\psi)_+ = (x_{ij} \mid i < r)R.$$

Using the inductive hypothesis, it remains to verify that  $a(T) < \mathbf{0}$ . By condition (1), for all integers  $1 \leq j \leq r - 1$ , we have

$$[H_{\mathfrak{M}}^i(R)^{\varphi_j}]_{\geq 0} = 0 \quad \text{for all } i \geq 0,$$

and using Lemma 5.5 it follows that

$$[H_{\mathfrak{p}+\mathfrak{b}}^i(R)^{\varphi_j}]_{\geq 0} = 0 \quad \text{for all } i \geq 0.$$

Consequently  $a(T^{\varphi_j}) < 0$  for  $1 \leq j \leq r - 1$ , which completes the proof.  $\square$

### 6. F-regularity

For the theory of tight closure, we refer to the papers [HH1,HH2] and [HH3]. We summarize results about F-rational and F-regular rings:

**Theorem 6.1.** *The following hold for rings of prime characteristic.*

- (1) *Regular rings are F-regular.*
- (2) *Direct summands of F-regular rings are F-regular.*
- (3) *F-rational rings are normal; an F-rational ring which is a homomorphic image of a Cohen–Macaulay ring is Cohen–Macaulay.*
- (4) *F-rational Gorenstein rings are F-regular.*
- (5) *Let  $R$  be an  $\mathbb{N}$ -graded ring which is finitely generated over a field  $R_0$ . If  $R$  is weakly F-regular, then it is F-regular.*

**Proof.** For (1) and (2) see [HH1, Theorem 4.6] and [HH1, Proposition 4.12] respectively; (3) is part of [HH2, Theorem 4.2], and for (4) see [HH2, Corollary 4.7], Lastly, (5) is [LS, Corollary 4.4].  $\square$



The characteristic zero aspects of tight closure are developed in [HH4]. Let  $K$  be a field of characteristic zero. A finitely generated  $K$ -algebra  $R = K[x_1, \dots, x_m]/\mathfrak{a}$  is of *F-regular type* if there exists a finitely generated  $\mathbb{Z}$ -algebra  $A \subseteq K$ , and a finitely generated free  $A$ -algebra

$$R_A = A[x_1, \dots, x_m]/\mathfrak{a}_A,$$

such that  $R \cong R_A \otimes_A K$  and, for all maximal ideals  $\mu$  in a Zariski dense subset of  $\text{Spec } A$ , the fiber rings  $R_A \otimes_A A/\mu$  are  $F$ -regular rings of characteristic  $p > 0$ . Similarly,  $R$  is of *F-rational type* if for a dense subset of  $\mu$ , the fiber rings  $R_A \otimes_A A/\mu$  are  $F$ -rational. Combining results from [Ha,HW,MS,Sm] one has:

**Theorem 6.2.** *Let  $R$  be a ring which is finitely generated over a field of characteristic zero. Then  $R$  has rational singularities if and only if it is of  $F$ -rational type. If  $R$  is  $\mathbb{Q}$ -Gorenstein, then it has log terminal singularities if and only if it is of  $F$ -regular type.*

**Proposition 6.3.** *Let  $K$  be a field of characteristic  $p > 0$ , and  $R$  an  $\mathbb{N}$ -graded normal ring which is finitely generated over  $R_0 = K$ . Let  $\omega$  denote the graded canonical module of  $R$ , and set  $d = \dim R$ .*

*Suppose  $R$  is  $F$ -regular. Then, for each integer  $k$ , there exists  $q = p^e$  such that*

$$\text{rank}_K R_k \leq \text{rank}_K [H_m^d(\omega^{(q)})]_k.$$

**Proof.** If  $d \leq 1$ , then  $R$  is regular and the assertion is elementary. Assume  $d \geq 2$ . Let  $\xi \in [H_m^d(\omega)]_0$  be an element which generates the socle of  $H_m^d(\omega)$ . Since the map  $\omega^{[q]} \rightarrow \omega^{(q)}$  is an isomorphism in codimension one,  $F^e(\xi)$  may be viewed as an element of  $H_m^d(\omega^{(q)})$  as in [Wa2].

Fix an integer  $k$ . For each  $e \in \mathbb{N}$ , set  $V_e$  to be the kernel of the vector space homomorphism

$$R_k \rightarrow [H_m^d(\omega^{(p^e)})]_k, \quad \text{where } c \mapsto cF^e(\xi). \tag{6.3.1}$$

If  $cF^{e+1}(\xi) = 0$ , then  $F(cF^e(\xi)) = c^p F^{e+1}(\xi) = 0$ ; since  $R$  is  $F$ -pure, it follows that  $cF^e(\xi) = 0$ . Consequently the vector spaces  $V_e$  form a descending sequence

$$V_1 \supseteq V_2 \supseteq V_3 \supseteq \dots$$

The hypothesis that  $R$  is  $F$ -regular implies  $\bigcap_e V_e = 0$ . Since each  $V_e$  has finite rank,  $V_e = 0$  for  $e \gg 0$ . Hence the homomorphism (6.3.1) is injective for  $e \gg 0$ .  $\square$

We next record tight closure properties of general  $\mathbb{N}$ -graded hypersurfaces. The results for  $F$ -purity are essentially worked out in [HR].

**Theorem 6.4.** *Let  $A = K[x_1, \dots, x_m]$  be a polynomial ring over a field  $K$  of positive characteristic. Let  $d$  be a nonnegative integer, and set  $M = \binom{d+m-1}{d} - 1$ . Consider the affine space  $\mathbb{A}_K^M$  parameterizing the degree  $d$  forms in  $A$  in which  $x_1^d$  occurs with coefficient 1.*

*Let  $U$  be the subset of  $\mathbb{A}_K^M$  corresponding to the forms  $f$  for which  $A/fA$  is  $F$ -pure. Then  $U$  is a Zariski open set, and it is nonempty if and only if  $d \leq m$ .*

Let  $V$  be the set corresponding to forms  $f$  for which  $A/fA$  is  $F$ -regular. Then  $V$  contains a nonempty Zariski open set if  $d < m$ , and is empty otherwise.

**Proof.** The set  $U$  is Zariski open by [HR, p. 156] and it is empty if  $d > m$  by [HR, Proposition 5.18]. If  $d \leq m$ , the square-free monomial  $x_1 \cdots x_d$  defines an  $F$ -pure hypersurface  $A/(x_1 \cdots x_d)$ . A linear change of variables yields the polynomial

$$f = x_1(x_1 + x_2) \cdots (x_1 + x_d)$$

in which  $x_1^d$  occurs with coefficient 1. Hence  $U$  is nonempty for  $d \leq m$ .

If  $d \geq m$ , then  $A/fA$  has  $a$ -invariant  $d - m \geq 0$  so  $A/fA$  is not  $F$ -regular. Suppose  $d < m$ . Consider the set  $W \subseteq \mathbb{A}_K^M$  parameterizing the forms  $f$  for which  $A/fA$  is  $F$ -pure and  $(A/fA)_{\bar{x}_1}$  is regular;  $W$  is a nonempty open subset of  $\mathbb{A}_K^M$ . Let  $f$  correspond to a point of  $W$ . The element  $\bar{x}_1 \in A/fA$  has a power which is a test element; since  $A/fA$  is  $F$ -pure, it follows that  $\bar{x}_1$  is a test element. Note that  $\bar{x}_2, \dots, \bar{x}_m$  is a homogeneous system of parameters for  $A/fA$  and that  $\bar{x}_1^{d-1}$  generates the socle modulo  $(\bar{x}_2, \dots, \bar{x}_m)$ . Hence the ring  $A/fA$  is  $F$ -regular if and only if there exists a power  $q$  of the prime characteristic  $p$  such that

$$x_1^{(d-1)q+1} \notin (x_2^q, \dots, x_m^q, f)A.$$

The set of such  $f$  corresponds to an open subset of  $W$ ; it remains to verify that this subset is nonempty. For this, consider

$$f = x_1^d + x_2 \cdots x_{d+1},$$

which corresponds to a point of  $W$ , and note that  $A/fA$  is  $F$ -regular since

$$x_1^{(d-1)p+1} \notin (x_2^p, \dots, x_m^p, f)A. \quad \square$$

These ideas carry over to multi-graded hypersurfaces; we restrict below to the bigraded case. The set of forms in  $K[x_1, \dots, x_m, y_1, \dots, y_n]$  of degree  $(d, e)$  in which  $x_1^d y_1^e$  occurs with coefficient 1 is parametrized by the affine space  $\mathbb{A}_K^N$  where  $N = \binom{d+m-1}{d} \binom{e+n-1}{e} - 1$ .

**Theorem 6.5.** Let  $B = K[x_1, \dots, x_m, y_1, \dots, y_n]$  be a polynomial ring over a field  $K$  of positive characteristic. Consider the  $\mathbb{N}^2$ -grading on  $B$  with  $\deg x_i = (1, 0)$  and  $\deg y_j = (0, 1)$ . Let  $d, e$  be nonnegative integers, and consider the affine space  $\mathbb{A}_K^N$  parameterizing forms of degree  $(d, e)$  in which  $x_1^d y_1^e$  occurs with coefficient 1.

Let  $U$  be the subset of  $\mathbb{A}_K^N$  corresponding to forms  $f$  for which  $B/fB$  is  $F$ -pure. Then  $U$  is a Zariski open set, and it is nonempty if and only if  $d \leq m$  and  $e \leq n$ .

Let  $V$  be the set corresponding to forms  $f$  for which  $B/fB$  is  $F$ -regular. Then  $V$  contains a nonempty Zariski open set if  $d < m$  and  $e < n$ , and is empty otherwise.

**Proof.** The argument for  $F$ -purity is similar to the proof of Theorem 6.4; if  $d \leq m$  and  $e \leq n$ , then the polynomial  $x_1 \cdots x_d y_1 \cdots y_e$  defines an  $F$ -pure hypersurface.

If  $B/fB$  is  $F$ -regular, then  $\mathfrak{a}(B/fB) < \mathbf{0}$  implies  $d < m$  and  $e < n$ . Conversely, if  $d < m$  and  $e < n$ , then there is a nonempty open set  $W$  corresponding to forms  $f$  for which the hypersurface

$B/fB$  is F-pure and  $(B/fB)_{\bar{x}_1\bar{y}_1}$  is regular. In this case,  $\bar{x}_1\bar{y}_1 \in B/fB$  is a test element. The socle modulo the parameter ideal  $(x_1 - y_1, x_2, \dots, x_m, y_2, \dots, y_n)B/fB$  is generated by  $\bar{x}_1^{d+e-1}$ , so  $B/fB$  is F-regular if and only if there exists a power  $q = p^e$  such that

$$x_1^{(d+e-1)q+1} \notin (x_1^q - y_1^q, x_2^q, \dots, x_m^q, y_2^q, \dots, y_n^q, f)B.$$

The subset of  $W$  corresponding to such  $f$  is open; it remains to verify that it is nonempty. For this, use  $f = x_1^d y_1^e + x_2 \cdots x_{d+1} y_2 \cdots y_{e+1}$ .  $\square$

## Acknowledgments

The authors would like to thank Shiro Goto and Ken-ichi Yoshida for their valuable comments.

## References

- [CHTV] A. Conca, J. Herzog, N.V. Trung, G. Valla, Diagonal subalgebras of bigraded algebras and embeddings of blow-ups of projective spaces, *Amer. J. Math.* 119 (1997) 859–901.
- [CH] S.D. Cutkosky, J. Herzog, Cohen–Macaulay coordinate rings of blowup schemes, *Comment. Math. Helv.* 72 (1997) 605–617.
- [Da1] V.I. Danilov, The group of ideal classes of a completed ring, *Math. USSR Sb.* 6 (1968) 493–500.
- [Da2] V.I. Danilov, Rings with a discrete group of divisor classes, *Math. USSR Sb.* 12 (1970) 368–386.
- [Fl] H. Flenner, Rationale quasihomogene Singularitäten, *Arch. Math.* 36 (1981) 35–44.
- [GG] A.V. Geramita, A. Gimigliano, Generators for the defining ideal of certain rational surfaces, *Duke Math. J.* 62 (1991) 61–83.
- [GGH] A.V. Geramita, A. Gimigliano, B. Harbourne, Projectively normal but superabundant embeddings of rational surfaces in projective space, *J. Algebra* 169 (1994) 791–804.
- [GGP] A.V. Geramita, A. Gimigliano, Y. Pitteloud, Graded Betti numbers of some embedded rational  $n$ -folds, *Math. Ann.* 301 (1995) 363–380.
- [GW1] S. Goto, K.-i. Watanabe, On graded rings. I, *J. Math. Soc. Japan* 30 (1978) 179–213.
- [GW2] S. Goto, K.-i. Watanabe, On graded rings. II  $\mathbb{Z}^n$ -graded rings, *Tokyo J. Math.* 1 (1978) 237–261.
- [Ha] N. Hara, A characterisation of rational singularities in terms of injectivity of Frobenius maps, *Amer. J. Math.* 120 (1998) 981–996.
- [HW] N. Hara, K.-i. Watanabe, F-regular and F-pure rings vs. log terminal and log canonical singularities, *J. Algebraic Geom.* 11 (2002) 363–392.
- [HH1] M. Hochster, C. Huneke, Tight closure, invariant theory, and the Briançon–Skoda theorem, *J. Amer. Math. Soc.* 3 (1990) 31–116.
- [HH2] M. Hochster, C. Huneke, F-regularity, test elements, and smooth base change, *Trans. Amer. Math. Soc.* 346 (1994) 1–62.
- [HH3] M. Hochster, C. Huneke, Tight closure of parameter ideals and splitting in module-finite extensions, *J. Algebraic Geom.* 3 (1994) 599–670.
- [HH4] M. Hochster, C. Huneke, Tight closure in equal characteristic zero, in preparation.
- [HHR] M. Herrmann, E. Hyry, J. Ribbe, On the Cohen–Macaulay and Gorenstein properties of multigraded Rees algebras, *Manuscripta Math.* 79 (1993) 343–377.
- [HIO] M. Herrmann, S. Ikeda, U. Orbanz, *Equimultiplicity and Blowing Up: An Algebraic Study*, Springer-Verlag, Berlin, New York, 1988.
- [HR] M. Hochster, J. Roberts, The purity of the Frobenius and local cohomology, *Adv. Math.* 21 (1976) 117–172.
- [Hy1] E. Hyry, Blow-up rings and rational singularities, *Manuscripta Math.* 98 (1999) 377–390.
- [Hy2] E. Hyry, The diagonal subring and the Cohen–Macaulay property of a multigraded ring, *Trans. Amer. Math. Soc.* 351 (1999) 2213–2232.
- [Lv1] O. Lavila-Vidal, On the Cohen–Macaulay property of diagonal subalgebras of the Rees algebra, *Manuscripta Math.* 95 (1998) 47–58.
- [Lv2] O. Lavila-Vidal, On the diagonals of a Rees algebra, thesis, Universitat de Barcelona, 1999.

- [LvZ] O. Lavila-Vidal, S. Zarzuela, On the Gorenstein property of the diagonals of the Rees algebra, *Collect. Math.* 49 (1998) 383–397.
- [LS] G. Lyubeznik, K.E. Smith, Strong and weak F-regularity are equivalent for graded rings, *Amer. J. Math.* 121 (1999) 1279–1290.
- [MS] V.B. Mehta, V. Srinivas, A characterization of rational singularities, *Asian J. Math.* 1 (1997) 249–271.
- [RSS] P. Roberts, A.K. Singh, V. Srinivas, Annihilators of local cohomology in characteristic zero, *Illinois J. Math.* 51 (2007) 237–254.
- [STV] A. Simis, N.V. Trung, G. Valla, The diagonal subalgebra of a blow-up algebra, *J. Pure Appl. Algebra* 125 (1998) 305–328.
- [Si] A.K. Singh, Veronese subrings and tight closure, *Pacific J. Math.* 192 (2000) 399–413.
- [Sm] K.E. Smith, F-rational rings have rational singularities, *Amer. J. Math.* 119 (1997) 159–180.
- [Tr] N.V. Trung, Diagonal subalgebras and blow-ups of projective spaces, *Vietnam J. Math.* 28 (2000) 1–15.
- [Wa1] K.-i. Watanabe, Rational singularities with  $k^*$ -action, in: *Commutative Algebra, Trento, 1981*, in: *Lect. Notes Pure Appl. Math.*, vol. 84, Dekker, New York, 1983, pp. 339–351.
- [Wa2] K.-i. Watanabe, F-regular and F-pure normal graded rings, *J. Pure Appl. Algebra* 71 (1991) 341–350.